

A Model Existence Theorem for Infinitary Formulas in Metric Spaces

Carlos E. Ortiz
Beaver College

February 1, 2008

Abstract

We prove a Model Existence Theorem for a fully infinitary logic L_A for metric structures. This result is based on a generalization of the notions of approximate formulas and approximate truth in normed structures introduced by Henson ([7]) and studied in different forms by Anderson ([1]) and Fajardo and Keisler ([2]). This theorem extends Henson's Compactness Theorem for approximate truth in normed space structures to infinitary formulas.

1 Introduction

In 1976 Henson in [6] introduced a logic of positive bounded formulas in Banach spaces to study the relationship between a Banach space E and its nonstandard hull $H(E)$. This logic L_{PB} is based on a first order language L containing a binary function symbol $+$, unary predicate symbols P and Q to be interpreted as the closed unit ball and the closure of its complement and, for every rational number r , a unary function symbol f_r to be interpreted as the operation of scalar multiplication by r . L_{PB} is closed under finite conjunction, disjunction and bounded quantification of the form $(\exists x)(P(x) \wedge \dots)$ or $(\forall x)(P(x) \Rightarrow \dots)$.

For any formula ϕ in L_{PB} and for every natural number n it is possible to define in a purely syntactical way a formula ϕ_n in L_{PB} , called the *n-approximation* of ϕ . Intuitively, ϕ_n is the formula that results from metrically

weakening the predicates that appear in ϕ in such a way that as n tends to ∞ , ϕ_n approaches ϕ . From this notion of approximations of formulas follows the definition of approximate truth: a formula $\phi \in L_{PB}$ is approximately true in a Banach space E (denoted by $E \models_{AP} \phi$) iff for every integer n , $E \models \phi_n$. It can be shown that $\models_{AP}$ is a weaker notion than $\models$.

The above definition of $\models_{AP}$ is the starting point of the model theory of Banach spaces. The cornerstone of the development of this model theory is Henson's Compactness Theorem ([10]). This theorem states that for particular collections of normed spaces models (we will call these collections "classes of models"), the following holds:

for any $\phi \in L_{PB}$, if every approximation of ϕ holds in some model of the class, then there is a model in the class where ϕ is approximately true and is also true.

Henson, Heinrich and Iovino developed the model theory of $\models_{AP}$ in normed spaces ([7, 8, 9, 5, 10]). It follows from their work that $\models_{AP}$ is the natural semantic notion to do model theory in normed structures since many of the usual theorems of Model Theory (Compactness, Lowenheim-Skolem, etc.) hold for normed structures (using the semantic notion of $\models_{AP}$). This is not the case for the usual semantic notion of $\models$ in normed structures (see [11]). Furthermore, $\models_{AP}$ is a concept that appears naturally in analysis. For example, "The continuous map $T : E \rightarrow E$ has a fixed point" is approximately true in E iff T has an almost fixed point ([3]).

However, there is a fundamental limitation to the above approach. Many of the relevant properties in functional analysis require a full infinitary logic to be expressed (e.g. reflexivity, superreflexivity). In order to extend the applicability of the tools and ideas of model theory to problems in functional analysis, it is desirable then to obtain model theoretic results for infinitary formulas in normed structures. A natural starting point is to obtain a Model Existence Theorem that generalizes Henson's Compactness Theorem to an infinitary logic and to general metric structures.

Furthermore, in areas like the geometry of Banach spaces, it is useful to have tools to construct spaces that satisfy complex infinitary properties. Famous problems solved recently, like the distortion problem ([18]) or the unconditional basis problem ([4]), require the construction of pathological Banach spaces. Problems still open, like the Fixed Point Property for reflexive spaces, would require (for a negative solution) the construction of

pathological Banach spaces with infinitary properties. Any model-theoretic tool used to construct metric spaces satisfying complex infinitary properties should be based on a Model Existence Theorem for infinitary logic in metric spaces.

In this paper we extend Henson's Compactness Theorem to a full infinitary logic (L_A) for metric structures. L_A is closed under countable conjunctions, negation and bounded existential quantification over countable many variables. This logic was introduced in [15] and studied extensively in [16]. The main tools used to obtain the Model Existence Theorem are the notions of approximate formulas and approximate truth for the formulas in L_A and the concept of a uniform collection of models.

As mentioned above, the logic L_A as well as the notion of approximate truth for L_A were introduced in [15]. For the sake of completeness, we define them here. The logic L_A is defined in Section 2. In this section we also define the analog of a collection of models in the classical first order logic: the uniform collection of models. In Section 3 we introduce a notion of approximate truth for the formulas in L_A that extends Henson's. The key idea here is to replace Henson's notion of a sequence of approximate formulas $\{\phi_n : n \in \omega\}$ (for formulas $\phi \in L_{PB}$) by a tree of approximate formulas $\{\phi_{h,n} : h \in I(\phi) \ n \in \omega\}$ where each sequence $\phi_{h,1}, \phi_{h,2}, \dots, \phi_{h,n}, \dots$ is a branch of the tree. In this way a sentence ϕ is approximately true in a metric structure E iff there exists a branch of the tree of approximations of ϕ such that all the finitary formulas in this branch hold in E . For formulas in L_{PB} it can be shown that the branches of the tree of approximations are all logically equivalent. In this case, our notion of approximate truth coincides with Henson's original one.

In Section 4 we prove a Model Existence Theorem for L_A along the following lines:

If there exists a branch $\phi_{h,1}, \dots, \phi_{h,n}, \dots$ of the tree of approximate formulas of ϕ such that for every n there is a model $\mathbf{E}_n$ in the uniform collection of models $\mathcal{M}$ with $\mathbf{E}_n \models \phi_{h,n}$ then there exists a model $\mathbf{E}$ in $\mathcal{M}$ such that : $\mathbf{E} \models \phi$ and $\mathbf{E} \models_{AP} \phi$.

2 The Logic L_A

We will use the symbol $\mathfrak{R}$ to refer to the metric space of the real numbers with the usual metric. Likewise, the symbol Q denotes the rationals, and Q^+ the rationals strictly greater than zero.

We will use $\mathbf{E}$, $\mathbf{B}$ to denote metric models, letters $E, F, \dots$ to denote first order models and letters $A, X, Y, \dots$ to refer to sets. We reserve the letters f, g for function (or function symbols), and ϵ, δ for real numbers. Likewise, we use $m, n, r, \dots$ for integers.

We can visualize a metric language as being made of two pieces: a classical first order model and a metric part.

DEFINITION 2.1 *Definition of a Metric Language*

Fix a first order language L made of function, relation and constant symbols (with finite arity). Fix also $M_{\mathfrak{R}}$ a collection of function symbols disjoint from the function symbols in L . Let $D_{\mathfrak{R}}$ be the collection of all the compact sets in $\mathfrak{R}$ with the usual metric.

$L(M_{\mathfrak{R}}, D_{\mathfrak{R}})$, the metric language induced by L , $M_{\mathfrak{R}}$ and $D_{\mathfrak{R}}$, is the collection of all the relation, function and constants symbols appearing in $L \cup M_{\mathfrak{R}}$, as well as the collection of all the compact sets in $\mathfrak{R}$.

The function symbols in $M_{\mathfrak{R}}$ are called the real valued function symbols of the language. These function symbols are going to be interpreted as maps taking values in $\mathfrak{R}$. There is a fixed real valued function symbol $\rho \in M_{\mathfrak{R}}$ with arity two. This symbol will be interpreted in every metric model as a metric function. $\square$

EXAMPLE 2.2

A natural metric language to deal with normed structures is

$$L(\{\rho(x, y), ||x||\}, D_{\mathfrak{R}})$$

Here L is the first order language $L = (\{(x + y)\} \cup \{r(x) : r \in Q^+\}, \{B_q : q \in Q\}, \{0\})$ with $x + y$ a binary model valued function symbol, $r(x)$ (for $r \in Q$) a unary model valued function symbol, 0 a constant and B_q (for every positive rational q) a unary predicate (to be interpreted in the normed structures as the ball of radius q centered at the origin).

The metric part of the logic is made of all the compact subsets of $\mathfrak{R}$ (the set $D_{\mathfrak{R}}$), of the binary real valued function symbol $\rho(x, y)$ and of the unary real valued function symbol $||.||$.

We will call this metric language the language for normed structures. Note that every normed space can be seen in a natural way as a structure in this language. $\square$

We remark that this definition of metric language can be extended naturally to include, for any complete metric space (X, ρ) , (X, ρ) -valued function symbols, and the collection of the compact metric subsets of (X, ρ) . All the results proved in this paper extend naturally to this setting (see [16]). For simplicity's sake, we refrain of doing so here.

Note that the collection of the compact subsets of the reals ($D_{\mathfrak{R}}$) appears in every metric language. In order to reduce notation clutter we omit to mention this set in the descriptions of the metric languages. It is understood that every metric language contains this collection.

For the rest of this section we fix a metric language $L(M_{\mathfrak{R}})$.

We define now a logic L_A over the metric language $L(M_{\mathfrak{R}})$. The terms and the values of the terms are defined by induction.

DEFINITION 2.3 *Definition of terms in L_A*

1. Let $\{x_i | i \in \omega_1\}$ be a fixed collection of variables. The variables x_i are terms. Their arity is 1. They are considered model valued terms.
2. Given a collection of model valued terms $\{t_i : i \leq n\}$ and a real valued function symbol (or a model valued function symbol) f with arity n , $f(\vec{t}) = f(t_1, \dots, t_n)$ is a real valued term (or model valued term) with arity given by the cardinality of the collection of variables in $\{t_i : i \leq n\}$. $\square$

In summary, terms have finite arity and they can be model valued or real valued.

We are ready to define the formulas of the logic L_A by induction. As usual, for every formula ϕ , we will use the notation $\phi(\vec{x})$ to mean that the free variables of ϕ are among the components of the vector $\vec{x}$. Likewise, $\phi(\vec{x}_1, \vec{x}_2, \dots)$ means that the free variables of ϕ are among the components of the vectors $\vec{x}_1, \vec{x}_2, \dots$.

DEFINITION 2.4 *Atomic formulas in L_A*

An atomic formula in L_A would be any expression of one of the following forms:

1. $C(\vec{t})$, where C is a relation symbol in L and $\vec{t}$ is a vector of model valued terms such that its arity agrees with the arity of C .
2. $B(t)$, where $B \in D_{\mathbb{R}}$ is a compact subset of the reals and t is a real valued term. $\square$

DEFINITION 2.5 *Definition of formulas in L_A*

1. An atomic formula is a formula in L_A .
2. If $\phi_1, \phi_2, \dots \phi_i \dots (i < \omega)$ is a collection of formulas in L_A , then for every n ,

$$\bigwedge_{i=1}^n \phi_i \text{ and } \bigwedge_{i=1}^{\infty} \phi_i$$

are in L_A .

3. If ϕ is a formula in L_A then $\neg\phi$ is also a formula in L_A .
4. Consider a formula $\phi(\vec{y}_1, \dots \vec{y}_n, \dots, x)$ in L_A . Let $\vec{K} = (K_1, K_2, \dots)$ be a corresponding vector of relation symbols. The following formula is in L_A :

$$\exists(\vec{y}_1, \dots \vec{y}_n, \dots) \in (K_1, \dots K_n, \dots)(\phi(\vec{y}_1, \dots \vec{y}_n, \dots, \vec{x})). \square$$

We will abbreviate $\neg \wedge \neg$ by $\vee$ and $\neg(\phi \wedge \neg\psi)$ by $\phi \Rightarrow \psi$. To avoid very long formulas we will also abbreviate

$$\exists(y_1 \dots y_n \dots) \in (K_1, \dots K_n, \dots)(\phi(x_1, \dots x_n, \dots, \vec{y}))$$

by $\exists \vec{y} \in \vec{K} \phi(\vec{x}, \vec{y})$. Likewise $\neg \exists \vec{y} \in \vec{K} \neg \phi(\vec{y}, \vec{x})$ will be abbreviated by $\forall \vec{y} \in \vec{K} \phi(\vec{y}, \vec{x})$.

Finally, given a countable set A with a fixed enumeration $A = \{a_1, \dots, a_n, \dots\}$ and countable formulas $\{\phi_a\}_{a \in A}$ we understand by $\bigwedge_{a \in A} \phi_a$ the formula $\bigwedge_{n=1}^{\infty} \phi_{a_n}$.

Likewise, for any arbitrary integer r , we understand by $\bigwedge_{a \in A \uparrow r} \phi_a$ the formula $\bigwedge_{n=1}^r \phi_{a_n}$.

Notice that the formulas in L_A admit bounded quantification over infinitely many variables. This enhances the expressive power of L_A , as the next example shows.

EXAMPLE 2.6 *Expressive power of L_A*

The normed structure E is reflexive.

For any Banach space $(X, \|\cdot\|)$, let B_1 denote the unitary ball. A characterization of reflexivity due to James ([12]) (see also [17]) that does not require any mention of the dual is the following:

A Banach space $(X, \|\cdot\|)$ is reflexive iff

$$\forall \epsilon > 0 \forall \{x_i\}_{i=1}^\infty \subseteq B_1 \exists k \in \omega \text{ dist}[\text{conv}(\{x_1, \dots, x_k\}), \text{conv}(\{x_{k+1}, \dots\})] \leq \epsilon$$

Here, for any set $A \subset X$, $\text{conv}(A)$ is the convex hull spawned by A . Similarly, given two sets $A, B \in X$, $\text{dist}[A, B] = \inf \{\|x - y\| : x \in A, y \in B\}$.

Recall the language for normed structures discussed in Example 2.2, $L = (\{\rho(x, y), \|\cdot\|\})$, with L the first order language

$$L = (\{(x + y)\} \cup \{r(x) : r \in Q^+\}, \{B_q : q \in Q^+\}, \{0\})$$

The following sentence of L_A expresses reflexivity in this language:

$$\bigwedge_{n=1}^\infty \forall (x_1, \dots, x_m, \dots) \in (B_1, B_1, \dots, B_1, \dots) \\ \bigvee_{k=1}^\infty \bigvee_{r=1}^\infty \bigvee_{\vec{a} \in CO(k)} \bigvee_{\vec{b} \in CO(r)} \left\| \sum_{i=1}^k a_i x_i - \sum_{j=1}^r b_j x_{k+j} \right\| \leq (1/n)$$

Here, $\forall s \in \omega$, $CO(s)$ is the subset of Q^s made of all the s -tuples $(a_1, \dots, a_s)$ such that $\sum_{i=1}^s a_i = 1$ and $a_1, \dots, a_s \geq 0$.

The normed structure E is uniformly convex. A normed space $(X, \|\cdot\|)$ is uniformly convex iff $\forall \epsilon > 0 \exists \delta > 0$ such that for every x, y in the unitary sphere, $\|x - y\| \geq \epsilon \Rightarrow \frac{\|x+y\|}{2} \leq 1 - \delta$.

We use again the language for normed structures $L(\{\rho, \|\cdot\|\})$. Here is how we express uniform convexity by a sentence in L_A :

$$\bigwedge_{n=1}^\infty \bigvee_{m=1}^\infty \forall (x, y) \in (B_1, B_1) [\|x - y\| \geq (1/n) \Rightarrow (1/2)\|x + y\| \leq (1 - 1/m)] \square$$

2.1 Semantics for L_A

The definition of the structures of L_A is a natural generalization of Henson's notion of a Normed Space Structure (see[10]).

Any model of a metric language $L(M_{\mathbb{R}})$ can be imagined as a classical first order model plus interpretations for the metric part of $M_{\mathbb{R}}$.

DEFINITION 2.7 *Definition of Model*

A model for $L(M_{\mathbb{R}})$ is a pair $\mathbf{E} = (E, F_{\mathbb{R}})$ where:

1. E is a model of the first order language L with interpretations for the function, relation and constant symbols in L . Let us call X the universe of E .
2. the real valued function symbol ρ is interpreted as a metric in X . Furthermore, every interpretation of a model valued function symbol $f \in L$ is a continuous function $f^{\mathbf{E}}$ (with respect to the product topology induced by ρ in X^{a_f}) from X^{a_f} to (X, ρ) .
3. $F_{\mathbb{R}} = \{f^{\mathbf{E}} \mid f \in M_{\mathbb{R}}\}$ with the property that for every real valued $f \in M_{\mathbb{R}}$ with arity a_f , the interpretation $f^{\mathbf{E}}$ is a continuous function, with respect to the product topology induced by ρ in X^{a_f} , from X^{a_f} to $\mathbb{R}$. $\square$

Once the interpretations for the symbols of $L(M_{\mathbb{R}})$ are defined one can define the interpretations of the terms in the model, $t^{\mathbf{E}}$, in the natural way. When the model and the interpretations of the elements of the language are clear from the context we will drop the symbol $t^{\mathbf{E}}$ and use t .

The definition of satisfaction follows in the standard way.

DEFINITION 2.8

The truth ($\models$) relation in the model $\mathbf{E} = (E, F_{\mathbb{R}})$ is constructed by induction in formulas in the usual way from the truth definition for the atomic case:

- Let $K(\vec{t}(\vec{x}))$ be an atomic formula in L_A with $K \in L$. Let $\vec{a} \in X^{|\vec{x}|}$. $\mathbf{E} \models K(\vec{t}(\vec{a}))$ iff $E \models K(\vec{t}(\vec{a}))$ in the usual first order sense.
- Let $B(t(\vec{x}))$ be an atomic formula with B a compact subset of the reals ($B \in D_{\mathbb{R}}$). Let $\vec{a} \in X^{|\vec{x}|}$. $\mathbf{E} \models B(t(\vec{a}))$ iff it is true that $t^{\mathbf{E}}(\vec{a}) \in B$. $\square$

2.2 Uniform Collection of Metric Models

The analog to the notion of collection of models for a first order language is given by the notion of uniform collection of models in metric languages.

Intuitively, a uniform collection of models is a collection of all models for $L(M_{\mathbb{R}})$ satisfying the same uniform continuity requirements for the interpretations of the function symbols and satisfying the same uniform “bound” for the interpretations of the predicate symbols. Recall that in this section we are dealing with a fix metric language $L(M_{\mathbb{R}})$.

DEFINITION 2.9 *Uniform Collection of Models*

Fix the following assignments:

- For every model valued (real valued) function symbol f with arity r , for every vector $\vec{K} = (K_1, \dots, K_r)$ of predicate symbols in L , fix a predicate symbol $K^{(f, \vec{K})}$ in L (or a compact predicate $D^{(f, \vec{K})}$ in $D_{\mathbb{R}}$). It is intended for $K^{(f, \vec{K})}$ ($D^{(f, \vec{K})}$) to contain the set $f(\vec{K})$ in every model of the class.
- For every two relation symbols Q, A with arities a, b in L , fix a predicate symbol $K^{(Q, A)}$ in L . It is intended for $K^{(Q, A)}$ to be a uniform cover of the union of the projections of the relations Q, A into the model.
- For every model valued (or real valued) function symbol f with arity r , for every vector $\vec{K} = (K_1, \dots, K_r)$ of predicate symbols in L , for every rational $\epsilon > 0$ fix a rational $\delta^{(f, \vec{K}, \epsilon)} > 0$.

A uniform collection of models for the above assignments is the collection of all the models $\mathbf{E} = (E, F_{\mathbb{R}})$ in $L(M_{\mathbb{R}})$ that satisfies:

1. **Uniform bound for function symbols.** For every model valued (real valued) function symbol f in L , for every vector $\vec{K} = (K_1, \dots, K_r)$ of predicate symbols in L ,

$$\mathbf{E} \models \forall \vec{x} \in \vec{K} \ K^{(f, \vec{K})}(f(\vec{x}))$$

(or

$$\mathbf{E} \models \forall \vec{x} \in \vec{K} \ D^{(f, \vec{K})}(f(\vec{x})))$$

2. **The relation symbols of L form a directed set.** For every relation symbols Q, A in L with arities a, b ,

$$\mathbf{E} \models \forall \vec{x}, \vec{y} \in (Q, A) \left(\bigwedge_{i=1}^a K^{(Q,A)}(x_i) \wedge \bigwedge_{j=1}^b K^{(Q,A)}(y_j) \right)$$

Furthermore $\mathbf{E}$ verifies that for every a in $\mathbf{E}$ there exists a predicate symbol K in L such that $(\mathbf{E}, F_{\mathfrak{R}}) \models K(a)$.

3. **Uniform continuity for function symbols.** For every model valued (or real valued) function symbol f with arity a , for every vector $\vec{K} = (K_1, \dots, K_a)$ of predicate symbols in L , for every rational $\epsilon > 0$,

$$\mathbf{E} \models \forall \vec{x}, \vec{y} \in (\vec{K}, \vec{K}) \left(\bigwedge_{i \leq a} \rho(x_i, y_i) < \delta^{(f, \vec{K}, \epsilon)} \Rightarrow \rho(f(\vec{x}), f(\vec{y})) \leq \epsilon \right)$$

(or

$$\mathbf{E} \models \forall \vec{x}, \vec{y} \in (\vec{K}, \vec{K}) \left(\bigwedge_{i \leq a} |x_i - y_i| < \delta^{(f, \vec{K}, \epsilon)} \Rightarrow \rho_Y(f(\vec{x}), f(\vec{y})) \leq \epsilon \right)) \square$$

EXAMPLE 2.10 *Uniform Class of Normed Space Structures*

Consider the metric language $L(\{\rho(x, y), ||x||\})$ with L the first order language $L = (\{(x + y)\} \cup \{r(x) : r \in Q\}, \{B_q : q \in Q^+\}, \{0\}, \}$ and with $\{\rho(x, y), ||x||\}$ as discussed in Example 2.2.

The natural uniform collection of models associated with this language and with the normed spaces is spawned by the following natural assignments:

- For $\rho(x, y)$ and for every B_{q_1}, B_{q_2} , let

$$D^{(\rho, (B_{q_1}, B_{q_2}))} = \{z \in \mathfrak{R} : 0 \leq z \leq q_1 + q_2\}.$$

For $(x + y)$ and for every B_{q_1}, B_{q_2} , let

$$K^{(+, (B_{q_1}, B_{q_2}))} = B_{q_1 + q_2}.$$

For $||x||$ and B_q let $D^{(||\cdot||, B_q)} = \{z \in \mathfrak{R} : 0 \leq z \leq q\}.$

- For every model valued function $r(x)$, with $r \in Q$, and for every B_q we assign $K^{(r(\cdot), B_q)} = B_{|r|q}$. For every B_{q_1}, B_{q_2} , we assign $K^{(B_{q_1}, B_{q_2})} = B_{q_1+q_2}$.
- For $\rho(x, y)$, for every B_{q_1}, B_{q_2} and for every rational $\epsilon > 0$, let $\delta^{(\rho, (B_{q_1}, B_{q_2}), \epsilon)} = \epsilon/2$. For $(x + y)$, for every B_{q_1}, B_{q_2} and for every rational $\epsilon > 0$, let $\delta^{(+, (B_{q_1}, B_{q_2}), \epsilon)} = \epsilon/2$. For $\|x\|$, B_q and a rational $\epsilon > 0$, let $\delta^{(\|\cdot\|, B_q, \epsilon)} = \epsilon$. For every function $r(x)$, with $r \in Q$, for every B_q and every rational $\epsilon > 0$, we assign $\delta^{(r(\cdot), B_q)} = \epsilon/r$.

It is clear that every model $\mathbf{E} = (E, \{\rho, \|\cdot\|\})$ of $L(M_{\mathbb{R}})$ such that $E = (X, \{0\}, +, \{r(x) : r \in Q\})$ is a vector space (over Q) and $\|\cdot\|$ is a norm in X , belongs to the above uniform collection of models. $\square$

3 Approximate Formulas in L_A

We define a notion of approximate formula (for the formulas in L_A) that extends Henson's definition (see [10]).

Our intention is to generate approximations of all the formulas in L_A by using the finitary formulas in L_A as building blocks. Inspired by the Souslin operation used in classical descriptive set theory (see for example [13]) to generate the analytic sets from the closed sets in a Polish space, we associate to every formula ϕ in L_A a tree of finitary formulas in L_A .

Formally, we associate to every formula ϕ in L_A a set of indices $I(\phi)$ (the branches of the tree of approximate formulas) and a set $\{\phi_{h,n} | h \in I(\phi), n \in \omega\}$ of finitary formulas in L_A . Intuitively, for every branch $h \in I(\phi)$, the approximate formulas in the branch (the collection $\{\phi_{h,n} | n \in \omega\}$) are going to "approach" ϕ as n tends to ∞ .

The notions of $I(\phi)$ and $\phi_{h,n}$ were introduced in [15] and studied in detail in [16].

Notation: sometimes, to emphasize that the pair h, n affects the formula ϕ , we will write $(\phi)_{h,n}$ for the approximation $\phi_{h,n}$.

Recall that we are working with a fixed metric language $L(M_{\mathbb{R}})$. Given two formulas ϕ, σ , we will write $\phi \equiv \sigma$ if ϕ and σ are identical formulas.

DEFINITION 3.1 *Definition of approximate formulas in L_A*

For any formula $\phi(\vec{x})$ in L_A we define:

1. A set of branches $I(\phi)$,
2. For any $h \in I(\phi)$ and $n \in \omega$, a finitary formula $\phi_{h,n}(\vec{x}) \in L_A$.

The definition is by induction in formulas as follows:

Atomic. For any atomic formula $C(t_1, \dots, t_r)$ or $D(t)$:

- The set $I(C(\vec{t})) = I(D(\vec{t})) = \{\emptyset\}$
- If $C \in L$, then for every h in $I(C(\vec{t}))$, for every integer n , $(C(\vec{t}))_{h,n} \equiv \exists \vec{y} \in C \bigwedge_{i=1}^r \rho(t_i, y_i) \leq (1/n)$ (this is the $1/n$ metric deformation of C).
- If $B \in \mathcal{M}_{\mathfrak{R}}$ is a compact subset of the reals, then the set $B_n = \{y \in \mathfrak{R} \mid \exists x \in B \text{ with } |x - y| \leq (1/n)\}$ is also a compact subset of $\mathfrak{R}$. We define then $(B(t))_{h,n} = B_n(t)$ (this is the $(1/n)$ deformation of the compact set B).

Conjunction. For any countable collection $\{\phi_i\}_{i=1}^\infty$ of formulas in L_A , we define:

- $I(\bigwedge_{i=1}^\infty \phi_i(\vec{x})) = \prod_{i=1}^\infty I(\phi_i)$ (the cartesian product of the $I(\phi_i)$).
- For every h in $I(\bigwedge_{i=1}^\infty \phi_i)$, for every integer n ,

$$\left(\bigwedge_{i=1}^\infty \phi_i \right)_{h,n} \equiv \bigwedge_{i=1}^n (\phi_i)_{h(i),n}$$

Negation. For any formula ϕ in L_A , we have:

- $I(\neg\phi) \subseteq (I(\phi) \times \omega)^\omega$ is the collection of all maps $f = (f_1, f_2)$ with the following “weak” surjectivity property:

$$\forall h \in I(\phi) \exists s \in \omega, (\phi)_{h, f_2(s)} \equiv (\phi)_{f_1(s), f_2(s)}$$

- For every $h \in I(\neg\phi(\vec{x}))$, for every integer n , $(\neg\phi)_{h,n} \equiv \bigwedge_{i=1}^n \neg (\phi_{h_1(i), h_2(i)})$

Existential. For every formula $\phi((\vec{v}_1, \vec{v}_2, \dots, \vec{v}_i, \dots), \vec{x})$, for every vector of bounding sets $\vec{K} = (K_1, K_2, \dots, K_i, \dots)$ of corresponding arities, we have:

- $I(\exists \vec{v} \in \vec{K} \phi(\vec{v}, \vec{x})) = I(\phi(\vec{v}, \vec{x}))$
- For every h in $I(\exists \vec{v} \in \vec{K} \phi(\vec{v}, \vec{x}))$, for every integer n , we have that

$$(\exists \vec{v} \in \vec{K} \phi(\vec{v}, \vec{x}))_{h,n} \equiv \exists \vec{v} \in \vec{K}(\phi(\vec{v}, \vec{x}))_{h,n}. \quad \square$$

We introduce some notation.

We call $I(\phi)$ **the set of branches** of ϕ . The formulas $\phi_{h,n}$ are the **approximate formulas** of ϕ , and the collection $T(\phi) = \{\phi_{h,n} : h \in I(\phi), n \in \omega\}$ is the **approximation tree** of the formula ϕ .

DEFINITION 3.2 *Approximate Truth*

Fix a model $\mathbf{E}$. Let $\phi(\vec{x})$ be an arbitrary formula in L_A . Let $\vec{b}$ be a vector of elements of X . We say that

$$\mathbf{E} \models_{AP} \phi(\vec{b}) \text{ (} \mathbf{E} \text{ approximately satisfy } \phi \text{)}$$

iff $\exists h \in I(\phi(\vec{x})) \forall n \in \omega, \mathbf{E} \models \phi_{h,n}(\vec{b})$.

Equivalently, $\mathbf{E} \models_{AP} \phi(\vec{b})$ iff there is a branch h of the approximation tree of $\phi(\vec{x})$ such that for every integer n , $\mathbf{E} \models \phi_{h,n}$.

We say that $\mathbf{E} \models_{AP}^h \phi(\vec{b})$ iff for every integer n , $\mathbf{E} \models_{AP} \phi_{h,n}(\vec{b})$. $\square$

We illustrate these definitions with some examples.

EXAMPLE 3.3 *Isometric Inclusion*

Let $(Y, |\cdot|)$ be a separable normed space generated by the countable set of independent vectors $\{e_1, e_2, \dots, e_n, \dots\}$ of norm 1. It is easy to see that $(Y, |\cdot|)$ is isometrically embedded into a Banach space $(X, \|\cdot\|)$ iff there exist unitary vectors $v_1, v_2, \dots, v_n, \dots$ in X such that:

$$\bigwedge_{m=1}^{\infty} \bigwedge_{\vec{a} \in Q^m} \left| \left\| \sum_{i=1}^m a_i v_i \right\| - \left| \sum_{i=1}^m a_i e_i \right| \right| \leq 0$$

Consider the language for normed structures $L(\{\rho(x, y), \|x\|\})$ discussed in Example 2.2. Recall that any normed space can be seen as a structure of this language with the natural interpretations. By $\vec{B}_1$ we denote the vector: $(B_1, B_1, \dots)$.

We can express that $(Y, |\cdot|)$ is isometrically embedded in a model $\mathbf{E}$ by the following sentence in L_A :

$$\mathbf{E} \models \exists \vec{x} \in \vec{B}_1 \bigwedge_{m=1}^{\infty} \bigwedge_{\vec{a} \in Q^m} \left| \left\| \sum_{i=1}^m a_i x_i \right\| - \left| \sum_{i=1}^m a_i e_i \right| \right| \leq 0$$

On the other hand, the statement

$$\mathbf{E} \models_{AP} \exists \vec{x} \in \vec{B}_1 \bigwedge_{m=1}^{\infty} \bigwedge_{\vec{a} \in Q^m} \left| \left\| \sum_{i=1}^m a_i x_i \right\| - \left| \sum_{i=1}^m a_i e_i \right| \right| \leq 0$$

is equivalent to the statement (since the set of branches $I(\phi) = \{\emptyset\}$) that for every integer r

$$\mathbf{E} \models \exists \vec{x} \in \vec{B}_1 \bigwedge_{m=1}^r \bigwedge_{\vec{a} \in Q^m \uparrow r} \left| \left\| \sum_{i=1}^m a_i x_i \right\| - \left| \sum_{i=1}^m a_i e_i \right| \right| \leq 1/r$$

It follows easily that this, in turn, is equivalent to the fact that for every finite dimensional subspace G of Y and for every $\epsilon > 0$ there is a finite dimensional subspace H in $\mathbf{E}$ that is ϵ -isomorphic to G . This is exactly the definition of the concept of finite representability of a normed space Y into X (see [14]).

In summary: the approximate validity of the statement “ Y is isometrically embeddable in $\mathbf{E}$ ” is equivalent to the statement that Y is finitely representable in $\mathbf{E}$. It is well known that the two notions are not equivalent (see for example [14]). This shows that the concept of approximate truth is different from validity. $\square$

EXAMPLE 3.4

Consider a formula ϕ such that $I(\phi) = \{o\}$ has cardinality one. By definition $I(\neg\phi) = (\{o\} \times \omega)^\omega$. Clearly this set can be identified with $(\omega)^\omega$. Using the definition of approximate truth, we see that for every $h \in (\omega)^\omega$, for every integer n , $(\neg\phi)_{h,n} = \bigwedge_{s=1}^n \neg(\phi)_{o,h(s)}$.

It follows that for every structure $\mathbf{E}$, $\mathbf{E} \models_{AP} \neg\phi$ iff $\exists h \in (\omega)^\omega$ such that for every integer n , $\mathbf{E} \models \bigwedge_{s=1}^n \neg(\phi)_{o,h(s)}$. Clearly this is equivalent to say that there exists an integer r such that, $\mathbf{E} \models \neg(\phi)_{o,r}$. $\square$

EXAMPLE 3.5

Consider a countable collection of formulas $\{\phi^i : i \in A\}$ such that for every i , $I(\phi_i) = \{o_i\}$ has cardinality one. By definition, using Example 3.4, $I(\bigwedge_{i \in A} \neg \phi^i)$ can be identified with $(\omega)^{A \times \omega}$. Using the definition of approximate truth, we see that for every $H \in (\omega)^{A \times \omega}$, for every integer n , $(\bigwedge_{i \in A} \neg \phi^i)_{H,n} = \bigwedge_{i \in A \upharpoonright n} \bigwedge_{s=1}^n \neg (\phi^i)_{o_i, H(i,s)}$.

It follows that for every structure $\mathbf{E}$, $\mathbf{E} \models_{AP} \bigwedge_{i \in A} \neg \phi^i$ iff there exists $H : A \times \omega \rightarrow \omega$ such that for every integer n , $\mathbf{E} \models \bigwedge_{i \in A \upharpoonright n} \bigwedge_{s=1}^n \neg (\phi^i)_{o_i, H(i,s)}$. Clearly again, this is equivalent to say that for every $i \in A$ there exists an integer n_i such that $\mathbf{E} \models \neg (\phi^i)_{o_i, n_i}$. $\square$

REMARK 3.6

It is not difficult to see by induction in formulas that if the formula ϕ in L_A is a positive formula (i.e. a formula without negation) then $I(\phi) = \{\emptyset\}$ has only one element, i.e. there is only one branch in the approximation tree of ϕ . For formulas that are not positive, the set of paths of the approximation tree has cardinality bigger than one (2^ω).

Consider the collection $L_{PBA} \subseteq L_A$ of all the formulas containing the atomic formulas and closed under countable conjunction, finite disjunction, existential and universal bounded quantification. This collection clearly contains the Henson's logic L_{PB} . Furthermore, it can be shown directly (see [16]) that the branches of the tree of approximation for the formulas $\phi \in L_{PB}$ are all logically equivalent in the following sense: $\forall h, g \in I(\phi) \mathbf{E} \models_{AP}^h \phi$ iff $\mathbf{E} \models_{AP}^g \phi$. This means that the tree of approximations essentially contains only one branch h . It is not hard then to verify that for every formula in L_{PB} our notion of $\models_{AP}$ and Henson's notion of approximate truth coincide. $\square$

In the next lemma, we prove two basic properties of the tree of approximations of any formula in L_A . In particular we show that the number of approximate formulas $\phi_{h,n}$ for any formula ϕ is countable. We remark that this lemma was already proved in [15]. We include the proof here for the sake of completeness.

LEMMA 3.7

For every formula $\phi(\vec{x}) \in L_A$:

1. For any model $\mathbf{E}$, for any vector of elements in $\vec{b}$ in $\mathbf{E}$, for any integer n and for all $h \in I(\phi)$,

$$\mathbf{E} \models \phi_{h,n+1}(\vec{b}) \Rightarrow \phi_{h,n}(\vec{b})$$

2. For every formula ϕ there exists a set $Dense(\phi) \subseteq I(\phi)$ at most countable with the following property:

For every branch h in $I(\phi)$, for every integer n , there exists a branch g in $Dense(\phi)$ such that: $\phi_{h,n}(\vec{x}) \equiv \phi_{g,n}(\vec{x})$ $\square$

PROOF:

1) By induction on the complexity of the formulas and Definition 3.1 of approximate formulas. Left to the reader.

2) By induction on the complexity of the formulas in L_A .

Atomic Formulas. Let $C(\vec{t})$ (or $B(t)$) be an atomic formula. Then we know that $I(C(\vec{t})) = \{\emptyset\}$ (or $I(B(t)) = \{\emptyset\}$) and $\forall n \in \omega \forall h \in I(C(\vec{t}))$ (or in $B(t)$):

$$(C(\vec{t}))_{h,n} = C_n(\vec{t}) \text{ (or } (B(t))_{h,n} = B_n(t))$$

Define then $Dense(C(\vec{t})) = I(C(\vec{t})) = \{\emptyset\}$ (or $Dense(B(t)) = I(B(t)) = \{\emptyset\}$). It is easy to see that this set verifies the desired property.

For the connectives and quantifier steps let us assume as induction hypothesis that for formulas ψ of less complexity than the formula ϕ , $I(\psi)$ and $Dense(\psi)$ satisfy the desired properties.

Conjunction. Consider the formula $\phi(\vec{x}) = \bigwedge_{i=1}^{\infty} \phi_i(\vec{x})$.

Recall that $I(\phi) = \prod_{i=1}^{\infty} I(\phi_i)$ and that for all integers n and for all $h \in I(\phi)$, $(\phi(\vec{x}))_{h,n} = \bigwedge_{i=1}^n (\phi_i(\vec{x}))_{h(i),n}$.

The construction of $Dense(\bigwedge_{i=1}^{\infty} \phi_i)$ is as follows:

$\forall n \in \omega$, let

$$p_n : \prod_{i=1}^{\infty} Dense(\phi_i) \mapsto \prod_{i=1}^n Dense(\phi_i)$$

be the usual projection map. By the induction hypothesis, $\forall i \in \omega$, $Dense(\phi_i)$ is at most countable. It is possible then to find, for every integer n , countable sets $Q_n \subseteq \prod_{i=1}^{\infty} Dense(\phi_i) \subset I(\bigwedge_{i=1}^{\infty} \phi_i)$ such that:

$$p_n(Q_n) = \prod_{i=1}^n Dense(\phi_i)$$

We define $Dense(\bigwedge_{i=1}^{\infty} \phi_i) = \bigcup_{n=1}^{\infty} Q_n$.

Clearly $Dense(\bigwedge_{i=1}^{\infty} \phi_i) \subseteq I(\bigwedge_{i=1}^{\infty} \phi_i)$ and is at most countable. Furthermore, by the induction hypothesis, for every $h = (h_1, \dots, h_i, \dots) \in I(\bigwedge_{i=1}^{\infty} \phi_i)$, for every integer n , there exists $(g_1, \dots, g_n) \in \prod_{i=1}^n Dense(\phi_i)$ such that:

$$\bigwedge_{i=1}^n (\phi_i)_{h_i, n} \equiv \bigwedge_{i=1}^n (\phi_i)_{g_i, n}$$

It follows that for every integer n there exists $g \in Dense(\bigwedge_{i=1}^{\infty} \phi_i)$ such that:

$$(\bigwedge_{i=1}^{\infty} \phi_i)_{h, n} \equiv \bigwedge_{i=1}^n (\phi_i)_{h_i, n} \equiv \bigwedge_{i=1}^n (\phi_i)_{g_i, n} \equiv (\bigwedge_{i=1}^{\infty} \phi_i)_{g, n}$$

This is the desired result.

Negation. Consider the formula $\phi(\vec{x}) = \neg\psi(\vec{x})$ in L_A .

Recall that $I(\neg\psi) \subseteq (I(\psi) \times \omega)^\omega$ is the collection of all the functions $f = (f_1, f_2)$ with the property that $\forall h \in I(\psi) \exists s, \psi_{h, f_2(s)} \equiv \psi_{f_1(s), f_2(s)}$.

Recall also that $\forall n \in \omega \forall h = (h_1, h_2) \in I(\neg\psi) (\neg\psi)_{h, n} = \bigwedge_{i=1}^n \neg(\psi_{h_1(i), h_2(i)})$.

The construction of the set $Dense(\neg\psi)$ is as follows:

Let $A \subseteq I(\neg\psi)$ be the collection of all the functions $f = (f_1, f_2) \in I(\neg\psi)$ such that $Image(f_1) \subseteq Dense(\psi)$. From the induction hypothesis it follows that $\forall n \in \omega$ the set $(Dense(\psi) \times \omega)^{\{1, \dots, n\}}$ is at most countable. For any integer let

$$proj_n : A \mapsto (Dense(\psi) \times \omega)^{\{1, \dots, n\}}$$

be the natural projection map. It is possible to find for every integer n a countable set $O(n) \subseteq A \subseteq I(\neg\psi) \subset (I(\psi) \times \omega)^\omega$ such that

$$proj_n(O(n)) = proj_n(A)$$

Let then $Dense(\neg\psi) = \bigcup_{n=1}^{\infty} O(n)$.

Let us verify the desired property. First, it is clear that $Dense(\neg\psi)$ is at most countable. Fix now $h = (h_1, h_2) \in I(\neg\psi)$ and an integer n .

By induction hypothesis for every integer s there exists $g_s \in Dense(\psi)$ such that

$$\psi_{h_1(s), h_2(s)} \equiv \psi_{g_s, h_2(s)}$$

It follows that there exists $f \in A$ such that for every integer m

$$(\neg\psi)_{h, m} \equiv \bigwedge_{s=1}^m \neg(\psi)_{h_1(s), h_2(s)} \equiv \bigwedge_{s=1}^m \neg(\psi)_{f_1(s), f_2(s)} \equiv (\neg\psi)_{f, m}$$

Now, by definition of $O(n) \subseteq A$, there exists $g \in O(n)$ such that $g \upharpoonright \{1, \dots, n\} = f \upharpoonright \{1, \dots, n\}$. We obtain then

$$(\neg\psi)_{h,n} \equiv \bigwedge_{s=1}^n \neg (\psi)_{f_1(s), f_2(s)} \equiv \bigwedge_{s=1}^n \neg (\psi)_{g_1(s), g_2(s)} \equiv (\neg\psi)_{g,n}$$

and this is the desired result.

Existential. Fix a formula ϕ in L_A with free variables among the collection $\{\vec{v}_i | i < \omega\} \cup \{\vec{x}\}$. Let $\vec{K} = (K_1, K_2 \dots)$ a corresponding vector of predicate symbols with true sort. Consider the formula $\psi(\vec{x}) = \exists \vec{v} \in \vec{K} \phi(\vec{v}, \vec{x})$.

Recall: $I(\psi) = I(\phi)$. Also, recall that $\forall n \in \omega \forall h \in I(\psi(\vec{x}))$,

$$(\psi(\vec{x}))_{h,n} = \exists \vec{v} \in \vec{K} \phi_{h,n}(\vec{v}, \vec{x})$$

Define then $Dense(\psi) = Dense(\phi)$. The verification of the desired property is trivial. ■

A welcome consequence of this lemma is the following corollary that states that the sets $I(\phi)$ are non empty.

COROLLARY 3.8 *For every formula $\phi \in L_A$, $I(\phi)$ is non empty.* □

PROOF: By induction on formulas. The only interesting step is the negation step. Recall that $I(\neg\phi) \subseteq (I(\phi) \times \omega)^\omega$ is the collection of all maps $f = (f_1, f_2)$ such that:

$$\forall h \in I(\phi) \exists s \in \omega, \phi_{h, f_2(s)} \equiv \phi_{f_1(s), f_2(s)} \quad (1)$$

Lemma 3.7 states that $Dense(\phi)$ is countable. Hence the set $B = Dense(\phi) \times \omega \subseteq I(\phi) \times \omega$ is countable. Let $f : \omega \mapsto I(\phi) \times \omega$ be such that $Im(f) = B$. It is easy to check using Lemma 3.7 that f verifies statement 1 above, hence $f \in I(\neg\phi)$. ■

4 A Model Existence Theorem for L_A

Our intention is to prove a model existence theorem for uniform collections of models using an ultraproduct construction very similar to the ultraproduct construction for normed spaces (see for example [3]).

In this section we fix a metric language $L(M_{\mathfrak{R}})$.

Let us recall some notation. Let D a compact subset of the reals with the usual metric. Let $s = (s_1, s_2, \dots s_i \dots)$ ($i < \omega$) be a sequence of elements in D . It is well known that for every ultrafilter U over ω there exists a unique $x \in D$ so that:

$$\forall \epsilon > 0 \exists p \in U \forall i \in p |s_i - x| \leq \epsilon$$

We will denote the element x by $\lim_{i \in U} s_i$.

DEFINITION 4.1 *Ultraproduct Construction*

Fix a uniform collection of models $\mathcal{W}$. Consider a sequence of models $\{\mathbf{E}_i = (E_i, F_{\mathfrak{R}}^i) | i < \omega\}$ of $\mathcal{W}$. For every i , let X_i be the universe of the first order model E_i . For every f in $\mathcal{F}$ we will denote by f^i the interpretation of this function symbol on the model $\mathbf{E}_i$. Likewise for any relation symbol C in L , C^i is the interpretation of C in $\mathbf{E}_i$.

Recall that for every relation symbol $K \in L$, for every integer n , K_n is the $(1/n)$ metric deformation of K .

Define X to be the set:

$$\{g \in \prod_{i=1}^{\omega} X_i \mid \exists K \in L \text{ a predicate such that } , \forall n \in \omega \forall^* i \in \omega, \mathbf{E}_i \models K_n(g(i))\}$$

In other words, X is the collection of all the sequences g that eventually converge to some predicate K .

Given a regular ultrafilter $\mathcal{U}$ over ω , we define on X the usual equivalence relation: For arbitrary x, y in X , $x \sim_{\mathcal{U}} y$ iff for every $\epsilon > 0$ there exists $p \in \mathcal{U}$ such that $\forall i \in p \rho^i(x(i), y(i)) \leq \epsilon$. It is easy to verify that this is truly an equivalence relation. We will denote by $[x]$ the equivalence class of x .

The set $(X/\sim_{\mathcal{U}})$ is endowed with a metric ρ in the natural way: For every pair $[x], [y]$, find two representatives x, y such that there exist K^1, K^2 predicates satisfying for every integer i : $\mathbf{E}_i \models K^1(x(i)) \wedge K^2(y(i))$ (this is possible because $\mathcal{U}$ is regular).

By the property of directed sets of uniform collections of models we know that there exists a unary predicate K such that for every integer i , $\mathbf{E}_i \models K(x(i)) \wedge K(y(i))$. Invoking again the properties of the uniform collections of models, we get that there exists a compact set D in $\mathfrak{R}$ such that for every i , $\mathbf{E}_i \models \forall (z, v) \in (K, K) D(\rho^i(z, v))$. It makes sense then to define

$$\rho([x], [y]) = \lim_{i \in \mathcal{U}} \rho^i(x(i), y(i))$$

It follows from the definition of a uniform collection of models that the metric function is well defined.

Using the set $(X/\sim_{\mathcal{U}})$ we define the interpretations of L :

- For every model valued function symbol f in L with arity a we define $f : (X/\sim_{\mathcal{U}})^a \mapsto (X/\sim_{\mathcal{U}})$ by:

$$\forall [\vec{x}] \in (X/\sim_{\mathcal{U}})^a, f([\vec{x}]) = [(f^1(\vec{x}(1)), \dots, f^i(\vec{x}(i)), \dots)]$$

The properties of the uniform collection of models guarantee that the image of f is a subset of $(X/\sim_{\mathcal{U}})$ and that f is well defined.

- For every real valued function symbol f in $M_{\mathfrak{R}}$ with arity a we define $f : (X/\sim_{\mathcal{U}})^a \mapsto \mathfrak{R}$ by:

$$\forall [\vec{x}] \in (X/\sim_{\mathcal{U}})^a, f([\vec{x}]) = \lim_{i \in U} f^i(\vec{x}(i)).$$

The properties of the uniform collections of models once again guarantee that the images of f are well defined.

- For any relation symbol C in L with arity a we define the interpretation of C in $(X/\sim_{\mathcal{U}})^a$ as follows:

$$[\vec{x}] \in C \text{ iff for every integer } n, \exists p \in \mathcal{U} \text{ such that for every } i \in p, \mathbf{E}_i \models (C(\vec{x}(i)))_n$$

Such a structure is denoted by $\prod_{\mathcal{U}} \mathbf{E}_i$. $\square$

The following remark follows directly from the previous construction.

REMARK 4.2 *The ultraproduct construction yields a model of $L(M_{\mathfrak{R}})$.*

Let $\mathcal{W}$ be a uniform collection of models of L . Let

$$\{\mathbf{E}_i | i < \omega\}$$

be a sequence of models of $\mathcal{W}$. Let $\mathcal{U}$ be a regular ultrafilter over ω . Then the structure $\prod_{\mathcal{U}} \mathbf{E}_i$ is a model of L . $\square$

PROOF: Let us verify first that $\prod_{\mathcal{U}} \mathbf{E}_i$ is a model of L .

Clearly $((X/\sim_{\mathcal{U}}), d)$ is a metric space. Using the property of uniform continuity of the uniform collection of models $\mathcal{W}$ we obtain that the functions f are continuous. It is easy to verify that the interpretation of the relation symbols C in L with arity a are closed subsets of $(X/\sim_{\mathcal{U}})^a$ (in the product topology).

Likewise, it is easy to verify that the interpretations of the real valued functions symbols in $M_{\mathbb{R}}$ are continuous functions. ■

We introduce the following notation. Given any model $\mathbf{E}$, any $\vec{a} = (\vec{a}_1, \dots, \vec{a}_i, \dots)$ in $\mathbf{E}$ and any corresponding vector of true sort predicate symbols $\vec{K} = (\vec{K}_1, \dots, \vec{K}_i, \dots)$, by $\vec{K}(\vec{a})$ we understand $\bigwedge_{i=1}^{\infty} K_i(\vec{a}_i)$.

The next lemma states the basic relationship between approximate formulas of L_A and the ultraproduct.

LEMMA 4.3 *Property of the Ultraproduct*

Fix $\mathcal{W}$ a uniform collection of models in $L(M_{\mathbb{R}})$. Let

$$\{\mathbf{E}_i | i < \omega\}$$

be a sequence of models in $\mathcal{W}$. Fix $\mathcal{U}$ a regular ultrafilter over ω . For any formula $\phi(\vec{x}) \in L_A$, for any sequence $\{\vec{a}_r\}_{r=1}^{\infty}$ of vectors whose elements are in $(X/\sim_{\mathcal{U}})$, the following are equivalent:

- $\exists h \in I(\phi) \forall n \in \omega \forall^* r \in \omega \prod_{\mathcal{U}} \mathbf{E}_i \models \phi_{h,n}(\vec{a}_r)$
- $\exists h \in I(\phi)$ such that $\forall n \in \omega \forall^* r \in \omega \exists p \in \mathcal{U}$ satisfying:

$$\forall i \in p, \mathbf{E}_i \models \phi_{h,n}(\vec{a}_r(i)). \quad \square$$

PROOF: By induction on formulas.

Atomic Formulas. The proof follows directly from the definition of the interpretation of the predicates in the model $\prod_{\mathcal{U}} \mathbf{E}_i$, using the facts that the predicates with fixed sort are closed, the interpretation of the function symbols are continuous functions and that the number of branches in the tree of approximation of atomic formulas is 1.

Conjunction. Both directions are direct.

Negation. $\Rightarrow$. Suppose that $\exists g \in I(\neg\phi) \forall n \in \omega \forall^* r \in \omega \prod_{\mathcal{U}} \mathbf{E}_i \models (\neg\phi(\vec{a}_r))_{g,n}$. By the definition of the negation step for approximate formulas we get that for every h in $I(\phi(\vec{x}))$ there exists an integer n such that $\forall^* r \in \omega \prod_{\mathcal{U}} \mathbf{E}_i \models \neg\phi_{h,n}(\vec{a}_r)$.

Using the induction hypothesis on the formula ϕ , and the fact that $\mathcal{U}$ is an ultrafilter, we get that for all h in $I(\phi)$ $\exists n \forall^* r \in \omega$ there exists a p in $\mathcal{U}$ satisfying:

$$\forall i \in p \ \mathbf{E}_i \models \neg \phi_{h,n}(\vec{a}_r(i))$$

Recall that $Dense(\phi) \subset I(\phi)$ is a countable set of branches “dense” in $I(\phi)$ (see Lemma 3.7). Define then the set $W =$

$$\{(h, n) : h \in Dense(\phi) \wedge n \in \omega \wedge \forall^* r \in \omega \exists p \in \mathcal{U} \forall i \in p, \mathbf{E}_i \models \neg(\phi_{h,n}(\vec{a}_r(i)))\}$$

Since the above set is countable, it is easy to obtain a $g = (g_1, g_2) : \omega \mapsto Dense(\phi) \times \omega$ such that $Image(g) = W$. Using the properties of the dense countable set $Dense(\phi)$ (Lemma 3.7) it is a standard procedure to verify that this function is in $I(\neg\phi(\vec{x}))$.

Furthermore, by definition of W , $\forall n \forall^* r \in \omega$ there exists a p in $\mathcal{U}$ such that:

$$\mathbf{E}_i \models (\neg\phi(\vec{a}_r(i)))_{g,n} \equiv \bigwedge_{s=1}^n \neg(\phi(\vec{a}_r(i)))_{g_1(s), g_2(s)}$$

This is the desired result.

$\Leftarrow$. Similar to the previous proof. Left to the reader.

Existential. Here we use the full power of the induction hypothesis on arbitrary sequences $\{a_r\}_{r=1}^\infty$.

$\Rightarrow$. Suppose that $\exists h \in I(\phi) \forall n \forall^* r \in \omega$

$$\prod_{\mathcal{U}} \mathbf{E}_i \models (\exists \vec{x} \in \vec{K} \phi(\vec{a}_r, \vec{x}))_{h,n}$$

By the definition of approximate formulas, the above statement implies that $\exists h \in I(\phi) \forall n \forall^* r \in \omega$

$$\prod_{\mathcal{U}} \mathbf{E}_i \models \exists \vec{x} \in \vec{K}(\phi(\vec{a}_r, \vec{x}))_{h,n}$$

It follows that for every integer r there exists a vector

$$\vec{d}_r$$

such that $\forall n \forall^* r$

$$\prod_{\mathcal{U}} \mathbf{E}_i \models \vec{K}(\vec{d}_r) \wedge \phi_{h,n}(\vec{a}_r, \vec{d}_r)$$

Furthermore, by the definition of $\sim$ we can select the elements of $\vec{d}_r$ in such a way that for every integer i , $\mathbf{E}_i \models \vec{K}(\vec{d}_r(i))$.

Using the induction hypothesis, we obtain then that there exists $h \in I(\phi)$ such that $\forall n \forall^* r$ there exists $p \in \mathcal{U}$ with :

$$\forall i \in p \ \mathbf{E}_i \models (\vec{K})(\vec{d}_r(i)) \wedge \phi_{h,n}(\vec{a}_r(i), \vec{d}_r(i))$$

But this implies that $\forall n, \forall^* r, \exists p \in \mathcal{U}$,

$$\forall i \in p \ \mathbf{E}_i \models \exists \vec{x} \in \vec{K} \phi_{h,n}(\vec{a}_r(i), \vec{x})$$

This is the desired result.

$\Leftarrow$: Similar to the previous proof. Left to the reader.

This completes the proof of the lemma. ■

We use the previous result to show that the ultraproduct is a rich model, i.e. a model where $\models_{AP}$ and $\models$ coincide for all formulas in L_A .

THEOREM 4.4 *The Ultraproduct is Rich for L_A*

Fix a uniform collection of models $\mathcal{W}$ in $L(M_{\mathfrak{R}})$. Let $\{\mathbf{E}_i\}_{i=1}^\omega$ be a sequence of models in $\mathcal{W}$. Fix $\mathcal{U}$ a regular ultrafilter over ω . Then for every formula $\phi(\vec{x})$ in L_A and for every $\vec{a}$, a vector of elements in $\prod_{\mathcal{U}} \mathbf{E}_i$ with finite or infinite arity, the following are equivalent:

- $\prod_{\mathcal{U}} \mathbf{E}_i \models_{AP} \phi(\vec{a})$
- $\prod_{\mathcal{U}} \mathbf{E}_i \models \phi(\vec{a})$ □

PROOF: By induction in formulas. The atomic, negation and countable conjunction cases are direct and we leave them to the reader.

For the existential case, there is only one interesting direction. Suppose that $\prod_{\mathcal{U}} \mathbf{E}_i \models_{AP} \exists \vec{v} \in \vec{K} \phi(\vec{v}, \vec{a})$.

From Lemma 4.3 it follows that there exists $h \in I(\phi)$ such that for every integer $n \exists p \in \mathcal{U}$ such that:

$$\forall i \in p \ \mathbf{E}_i \models \exists \vec{v} \in \vec{K} \phi_{h,n}(\vec{v}, \vec{a}(i))$$

Since the ultrafilter is regular, it is easy to construct by diagonalization a sequence $\vec{b}$ of vectors of functions (elements of $\prod_{\mathcal{U}} \mathbf{E}_i$) corresponding to $\vec{v}$ and such that $\forall n \in \omega \exists p \in \mathcal{U} \forall i \in p$:

$$\mathbf{E}_i \models \vec{K}(\vec{b}(i)) \wedge \phi_{h,n}(\vec{b}(i), \vec{a}(i))$$

We again invoke Lemma 4.3 to obtain that $\prod_{\mathcal{U}} \mathbf{E}_i \models_{AP} \vec{K}(\vec{b}) \wedge \phi(\vec{b}, \vec{a})$.

By induction hypothesis we get $\prod_{\mathcal{U}} \mathbf{E}_i \models \vec{K}(\vec{b}) \wedge \phi(\vec{b}, \vec{a})$, and this implies $\prod_{\mathcal{U}} \mathbf{E}_i \models \exists \vec{v} \in \vec{K} \phi(\vec{v}, \vec{a})$. This completes the proof. ■

The main consequence of the previous lemmas is the following Model Existence Theorem.

THEOREM 4.5 *Model Existence Theorem for L_A*

Let $\mathcal{M}$ be a uniform collection of models for $L(M_{\mathfrak{R}})$. Fix a sentence $\phi \in L_A$. Suppose that there exists a branch $h \in I(\phi)$ such that for every integer n there exists $\mathbf{E}_n \in \mathcal{M}$ such that:

$$\mathbf{E}_n \models \phi_{h,n}$$

Then there exists a model $\mathbf{E}$ in $\mathcal{M}$ such that:

$$\mathbf{E} \models_{AP} \phi \text{ and } \mathbf{E} \models \phi \quad \square$$

PROOF: Select a regular ultrafilter over ω and apply Lemma 4.3 to the sequence of models $\{\mathbf{E}_n\}_{n=1}^{\infty}$. We obtain that $\prod_{\mathcal{U}} \mathbf{E}_i \models_{AP} \phi$.

Applying now Theorem 4.4 to $\prod_{\mathcal{U}} \mathbf{E}_i$ we get that $\prod_{\mathcal{U}} \mathbf{E}_i \models \phi$.

It remains to show that $\prod_{\mathcal{U}} \mathbf{E}_i$ is in the uniform collection $\mathcal{W}$. It is enough to check that $\prod_{\mathcal{U}} \mathbf{E}_i$ satisfies the same uniform bounds for the function symbols in $L(M_{\mathfrak{R}})$, the same inclusion relationship for the true sort predicates and the same uniform continuity property for the function symbols that the models in $\mathcal{W}$. Left to the reader. ■

REMARK 4.6

Recall that in Remark 3.6 we mentioned that for formulas $\phi \in L_{PB}$ the tree of approximations has only one branch h (up to logical equivalence), and that for this branch, the approximate formulas $\{\phi_{h,n}\}_{n=1}^{\infty}$ coincide with Henson's approximate formulas $\{\phi_n\}$.

It follows then that we have, as a corollary of the Model Existence Theorem, the following version of Henson's Compactness Theorem:

COROLLARY 4.7 *Henson's Compactness Theorem*

Fix $L(\{\rho, ||\cdot||\} \cup M_{\mathfrak{R}})$ with L and $\{\rho, ||\cdot||\}$ as in Example 2.2. Let $\mathcal{M}$ a uniform collection of normed structures in the above language. Fix a sentence $\phi \in L_{PB}$. If for every integer n there exists a model $\mathbf{E}_n \in \mathcal{M}$ with $\mathbf{E}_n \models \phi_n$, then there exists a model $\mathbf{E} \in \mathcal{M}$, such that $\mathbf{E} \models_{AP} \phi$ and $\mathbf{E} \models \phi$. □

References

- [1] Anderson R.M. “ “Almost” implies “Near”, ” *Trans. Amer. Math. Soc.* **296**, 1986, pp. 229-237.
- [2] Fajardo S., and H.J. Keisler H.J. “Neometric Forcing,” in preparation.
- [3] Aksoy A.G., and Khamsi M.A. *Nonstandard Methods in Fixed Point Theory*. New York: Springer Verlag, 1990.
- [4] Gowers W.T., and Maurey B. “The Unconditional Basic Sequence Problem,” *Journal A.M.S.* **6**, 1993, pp. 851-874.
- [5] Heinrich S., and Henson C.W. “Banach Space Model Theory,II: Isomorphic equivalence,” *Math. Nachr.* **125**, 1986, pp. 301-317.
- [6] Henson C.W. “When Do Two Banach Spaces Have Isometrically Isomorphic Nonstandard Hulls?,” *Israel J. Math.* **22**, 1975, pp. 57-67.
- [7] Henson C.W. “Nonstandard Hulls of Banach Spaces,” *Israel J. Math.* **25**, 1976, pp. 108-144.
- [8] Henson C.W. “Banach Space Model Theory III: Separable Spaces Under Isometric Equivalence,” in preparation.
- [9] Henson C.W. Forthcoming paper on model theory of L_p spaces.
- [10] Henson C.W., and Iovino J. “Banach Space Model Theory, I: Basics,” in preparation.
- [11] Iovino J. *Stable Theories in Functional Analysis*. Ph.D. thesis, University of Illinois at Urbana-Champaign, 1994.
- [12] James R.C. “Weak Compactness and Reflexivity,” *Israel J. Math.* **2**, 1964, pp 101-119.

- [13] Kechris A. S. *Classical Descriptive Set Theory*. New York: Springer Verlag, 1995.
- [14] Lindenstrauss J., and Tzafriri L. *Classical Banach Spaces*. Berlin: Springer Verlag, 1977.
- [15] Ortiz C.E. “Approximate Truth and Nonstandard Analysis,” *Cahiers du Centre de Logique* **9**. Louvain la Neuve, Academia Bruylant, 1996.
- [16] Ortiz C.E. *Truth and Approximate Truth in Metric Spaces*. Ph.D. thesis, University of Wisconsin-Madison, 1997.
- [17] Van Dulst D. *Reflexive and Superreflexive Banach Spaces*. Amsterdam: Mathematical Centre Tract, 1978.
- [18] Odell E., and Schlumprecht Th. “The Distortion Problem,” *Acta Math.* **173**, 1994, pp. 259-281.

Beaver College, Glenside, PA,
ortiz@beaver.edu

Mathematical Subject Classification, 1991:
Primary 03C65
Secondary 46B08, 46B20

Keywords: approximate truth, compactness theorem, normed space structures.